

Adoptive immunotherapy of cancer

Gene transfer of T cell specificity

Amir A. Al-Khami, Shikhar Mehrotra and Michael I. Nishimura*

Division of General Surgery; Department of Surgery; Medical University of South Carolina; Charleston, SC USA

Key words: T cell receptor, CAR, T cells, gene therapy, immunotherapy

Adoptive transfer of tumor-reactive T cells has emerged as a promising advance in tumor immunotherapy. Specifically, infusion of tumor-infiltrating lymphocytes has led to long-term objective clinical responses for patients with metastatic melanoma. Donor lymphocyte infusion is also an effective treatment of post-transplant lymphoproliferative disease. However, adoptive T cell therapy has restrictions in the isolation and expansion of antigen-specific lymphocytes for a large group of patients. One approach to circumvent this limitation and extend adoptive immunotherapy to other cancer types is the genetic modification of T cells with antigen-specific receptors. In this article, we review strategies to redirect T cell specificity, including T cell receptor gene transfer and antibody receptor gene transfer.

Adoptive Cell Transfer Therapy

Despite multiple approaches to therapy and prevention, cancer remains a major cause of death worldwide. Conventional therapies targeting dividing cells, using chemotherapy or radiation therapy, also affect normal cells and often fail in preventing the metastatic spread of the disease. Therefore, immunotherapy is an alternative modality of treatment that attempts to harness the specificity of the immune system to target tumor propagation without harming normal cells.¹ Active immunotherapy with therapeutic vaccines aims to elicit immune responses in vivo that will lead to an antitumor effect. With current cancer vaccine approaches, including tumor cell vaccines, peptide vaccines, viral vector vaccines, plasmid DNA vaccines and dendritic cell vaccines,² objective clinical responses have not yet been successfully achieved.³ In contrast, passive immunotherapeutic strategies with adoptive cell transfer (ACT) involve ex vivo stimulation of tumor-reactive T cells that are then transferred back to the patient.⁴

Adoptive Cell Transfer Following Allogeneic Bone Marrow Transplantation

The most successful T cell immunotherapy to date has possibly been the use of donor T cells following allogeneic hematopoietic

stem cell (HSC) transplantation. Indeed, treatment of patients with myeloid leukemias relapsing after allogeneic HSC transplantation by donor lymphocyte infusion has proved to induce long-term, complete remissions through graft-versus-leukemia reactivity. In this regard, the hematopoiesis-restricted minor histocompatibility antigens HA-1 or HA-2 expressed on malignant cells of the recipient can serve as target antigens for donor T cell recognition.^{5,6} The role of graft-versus-leukemia effect was underscored initially by observing complete hematologic and cytogenetic remission in 3 patients with relapsed chronic myeloid leukemia infused with donor buffy coat cells.⁷ Later clinical trials resulted in complete remissions in 70% of patients with chronic myeloid leukemia in response to donor lymphocyte transfusions. Complete remissions were also induced in 29% of patients with acute myeloid leukemia. In contrast, acute lymphoblastic leukemia did not respond to donor lymphocyte transfusions.⁸

During the immune recovery after HSC transplantation, reactivation of latent viruses, such as cytomegalovirus (CMV) and Epstein-Barr virus (EBV), is common and often causes an indicative disease.⁹ Yet, the adoptive transfer of T cell clones appeared effective in preventing complications of virus-associated infection and disease after transplant. For example, donor, CMV-specific CD8⁺ T cell clones were infused into recipients of the marrow transplant after stimulation with autologous fibroblasts infected with CMV. CMV-specific cytotoxic T lymphocytes (CTLs) were reconstituted and persisted in all patients.¹⁰ Besides, CMV-specific CTLs, stimulated with CMV lysate, were successfully used in the treatment of CMV infection not responding to prolonged antiviral chemotherapy.¹¹ The results of adoptive immunotherapy for the prophylaxis and treatment of EBV-associated, post-transplant lymphoproliferative disease (PTLD) were also efficacious. EBV-infected B cells of PTLD are highly immunogenic, expressing latent EBV antigens including the immunodominant EBV nuclear antigen-3.¹² Moreover, lymphoblastoid cell lines generated by infecting donor B cells with a laboratory strain of EBV have effectively been used as antigen-presenting cells for the expansion of EBV-specific CTLs in vitro.⁹ When donor, EBV-specific T cell lines were transferred into more than 60 marrow recipients as prophylaxis for PTLD, none of the 60 patients developed the malignancy. Of 6 additional patients that received virus-specific CTL after the onset of lymphoma, 5 patients underwent a complete regression.¹³

Overall, these studies demonstrating the efficiency of adoptive immunotherapy against allogeneic and viral antigens have

*Correspondence to: Michael I. Nishimura; Email: nishimur@musc.edu
Submitted: 04/14/11; Accepted: 04/14/11
<http://dx.doi.org/10.4161/self.2.2.15832>

profound implications for the development adoptive immunotherapy approaches to the treatment of patients with solid cancers.¹⁴

Adoptive Cell Transfer in Metastatic Melanoma

For patients with tumors that do not express viral antigens, the development of ACT therapies has had limited success, because the generation of tumor antigen-specific T cells has been problematic.¹⁵ The finding that normal lymphocytes, after culture with IL-2, caused a significant lysis of autologous tumor led to the use of these lymphokine activated killer (LAK) cells for the treatment of patients with metastatic cancers.¹⁶ Nevertheless, adoptive transfer of LAK cells plus IL-2 administration induced complete tumor regression in a limited number of patients.^{17,18} By applying advances in basic T cell biology and new techniques, better responses have been achieved later. The adoptive transfer of tumor-infiltrating lymphocytes (TILs) into patients with melanoma along with high-dose IL-2 resulted in an objective response rate of 34%. Most of these responses, however, were transient and patients had a limited persistence of the transferred cells.¹⁹ Another approach was to use CD8⁺ T cell clones specific for the melanoma antigens gp100 and MART-1 for the ACT therapy, in addition to low-dose or high-dose IL-2.^{20,21} Yet, T cell clones declined rapidly after adoptive transfer, and no objective clinical responses were reported. The clinical effectiveness of TIL therapy was further enhanced using nonmyeloablative, lymphodepleting chemotherapy in combination with ACT. In these protocols, patients with metastatic melanoma were infused with ex vivo-expanded, tumor-reactive TILs and high-dose IL-2 after a preconditioning regimen of cyclophosphamide and fludarabine.²² This approach resulted in a persistent clonal repopulation of T cells and objective tumor regression in 47% of patients. A following report on the same protocol showed an objective clinical response in 51% of patients.²³ More recently, up to 72% objective response rates were observed after increasing the intensity of the preparative regimens (12 Gy plus cyclophosphamide and fludarabine).²⁴

Despite encouraging results described above, the generation of tumor-reactive lymphocytes in this approach of immunotherapy is often a major limitation. Additionally, it is difficult to identify these tumor-reactive lymphocytes in many patients, in particular those with non-melanoma tumors. An attempt around this limitation is the genetic modification of T cells to redirect antigen specificity. The validity of this approach has been established by the long-term persistence of adoptively transferred, genetically modified T cells in humans.^{25,26} Furthermore, a uniform population of redirected antigen-specific cells can be rapidly generated by gene transfer in accessible peripheral blood T cells, in addition to TILs. The current approaches studied in this area involve: (1) naturally occurring two-chain TCR molecules and (2) single chain antibody constructs bound to intracellular T cell signaling domains.

T Cell Receptor Gene Transfer

The main determinant of tumor recognition by T cells is appropriate binding of the major histocompatibility complex

(MHC)-antigen complex by the TCR. Genes that encode the α and β chains are cloned from tumor-reactive T cells restricted to a particular HLA allele and then introduced into recipient T cells to endow them with the specificity of the donor TCR.²⁷ The first demonstration of redirected T cell specificity by TCR gene transfer was reported by Dembic et al. in 1986. With the identification of the TCR α and β chain sequences from two HLA-A2-restricted, MART-1/Melan A-reactive T cell clones,²⁹ Clay et al. were the first to report the transfer of TCR genes from one of these clones via a retroviral vector into human peripheral blood lymphocytes. Transduced cells acquired stable reactivity to MART-1/Melan A peptide and human HLA-A2 melanoma lines.^{30,31} Since these studies, the same approach was used to transfer the specificity for a number of tumor-associated antigens and viral antigens associated with tumor development.³²⁻³⁸

Several viral vectors have been tested for efficient transduction of human primary T lymphocytes. Adenoviruses were the first viral vectors to be used. Although adenoviral vectors can infect both dividing and nondividing cells, they are highly immunogenic, leading to strong (or preexisting) immune responses and rapid elimination of transduced cells *in vivo*.^{39,40} Gamma retroviral vectors were the first to provide efficient transduction. They have low immunogenicity, infect only proliferating cells and integrate into the host DNA, thus allowing stable transgene expression.⁴¹ Different classes of retroviruses that have been used for transfer of genes to human cells include lentiviral vectors. In contrast to retroviral vectors, lentiviruses can be used to infect both dividing and nondividing cells.^{42,43} Though, comparison of vectors is still lacking.

Mispairing of the introduced chains with endogenous TCR chains is a main concern in this approach; this may dilute the number of correctly expressed TCRs and could result in TCR heterodimers with unknown specificity, possibly inducing autoimmunity. Among those strategies developed to counter TCR mispairing are: (1) insertion of murine constant regions with the variable region of the human TCR⁴⁴ and (2) insertion of cysteine residues into the α and β chains that favors pairing of only the transduced chains.⁴⁵

One concern with TCR immunotherapy is the ligand-binding affinity of TCR transgenes. Therefore, selection of a TCR with high-affinity is crucial in improving the functional avidity of TCR-transduced cells. While isolating such a TCR from a patient is rare, this obstacle has recently been overcome by using methods such as phage display. This approach has produced TCRs with picomolar affinities, representing almost a million-fold improvement in affinity compared to the parent TCR.⁴⁶ High-affinity TCRs can also be obtained from HLA-A2 transgenic mice following immunization against human cancer antigens. This method usually results in high-avidity CTLs.⁴⁷

The first clinical example of this approach was reported by Morgan et al. at the NCI. Infusion of 15 patients with metastatic melanoma with their autologous lymphocytes transduced with genes encoding a TCR reactive against MART-1 following nonmyeloablative lymphodepletion resulted in a durable engraftment at levels exceeding 10% of peripheral blood lymphocytes for at least 2 months after treatment.⁴⁸ In this study, 2 patients (13%)

underwent objective regression of metastatic melanoma lesions. In another study, the same group attempted to use more reactive TCRs by: (1) screening highly reactive MART-1 T cell clones and (2) immunizing HLA-A2 transgenic mice with human gp100 peptide. Objective cancer regressions were seen in 30% and 19% of patients with metastatic melanoma, who received the human or mouse TCR, respectively. However, patients exhibited destruction of normal melanocytes in the skin, eye and ear, which was managed by steroid treatment.⁴⁹ More recently, autologous T lymphocytes genetically engineered to express a murine TCR against human carcinoembryonic antigen (CEA) were administered to 3 patients with metastatic colorectal cancer refractory to standard treatments. All patients showed decreased levels of serum CEA (74–99%), and one patient had an objective regression of cancer metastatic to the lung and liver. Though, a severe transient inflammatory colitis was induced in all three patients.⁵⁰

Chimeric Antibody Receptor Gene Transfer

Chimeric antibody receptors (CARs) use the extracellular single-chain variable fragment (scFv) of an antibody fused to intracellular signaling domains, such as the TCR ζ chain (CD3 ζ) or IgE high-affinity receptor (Fc ϵ RI γ).^{51,52} With such CARs, the recognition specificity of lymphocytes is redirected to desired cell surface tumor antigens, in a non-HLA restricted manner and without antigen processing and presentation by the target cell. Whereas a TCR binds only short peptides derived from protein antigens, CARs can recognize nonproteinous antigens, including carbohydrates and glycolipids. In addition, high-affinity scFv domain makes the transduced T cells more sensitive to low antigen densities.⁵³ However, a key to this approach is that the target antigens should be carefully chosen, so as to be highly expressed only on tumor cells and not on normal cells.⁵⁴

CARs have been developed to target antigenic molecules on various human tumors, including cancers of the breast, kidney, lung, colon, prostate, ovary, skin, pancreas and immune cells.^{54–60} Even though the first CARs were reported to redirect the cytotoxic function (tumor cell killing and cytokine production) of the engineered T cells, they did not transmit proliferative signals in response to antigen. To undergo full activation, T cells require signaling through their TCRs and costimulatory molecules, mainly CD28. As such, other intracellular signaling domains have been added to the CD3 ζ chain of CARs. Incorporating CD28 signaling domains resulted in enhanced proliferation, cytokine secretion, and tumor control.^{55,61,62} Inclusion of different costimulatory signaling domains, such as CD137 (4-1BB), CD134 (OX40) and ICOS (inducible T cell costimulator), has

been shown to further augment T cell survival and effector functions through CAR recognition.^{63–65}

The first CARs evaluated clinically were designed to target CD20 on lymphoma,⁶⁶ alpha-folate receptor on ovarian cancer,⁶⁷ carboxy-anhydrase-IX on renal cell carcinoma,⁵⁴ and the L1-cell adhesion molecule (CD171) and the diasialoganglioside GD2 on neuroblastoma.^{68,69} Overall, these studies were characterized by short-term persistence of gene-modified T cells and poor clinical responses. In the renal cell carcinoma trial, patients developed liver toxicity, probably due to the specific reactivity of T-cells against carboxy-anhydrase-IX-expressing bile duct epithelial cells.⁵⁴ Two adverse event reports, resulting in patient deaths, have recently been published.^{70,71} The first report targeted ERBB2 (HER-2/neu) in a patient with colon cancer metastatic to the lungs and liver.⁷⁰ Lymphodepletion followed by infusion of T cells transduced with a CAR vector containing CD28, 4-1BB, and CD3 ζ signaling domains caused immediate pulmonary toxicity. In this study, a cytokine storm has possibly been released by the administered cells in response to the low levels of ERBB2 expressed on normal lung epithelial cells. In the second report, chronic lymphocytic leukemia patients were treated with a CAR, comprising the CD28 cytoplasmic domain in addition to that of the CD3 ζ -chain.⁷¹ One patient, who received T cells following lymphodepleting chemotherapy, developed renal failure due primarily to hypotension as a consequence of sepsis-like syndrome.

Conclusion

Infusion of antigen-specific T cells, referred to as adoptive T cell therapy, has yielded therapeutic responses in patients with chronic infections and cancer. A better understanding of how human T cells function and modulate immune responses would largely increase the therapeutic potential of adoptive cell therapy. Recently, TCR or CAR gene transfer has been developed as a reliable method to generate ex vivo large numbers of T cells of a given antigen-specificity, for which TCRs or antibodies have been identified. The ability to develop modified TCR transgenes to prevent mispairing with endogenous TCR and generate CARs incorporating multiple costimulatory signaling domains could further improve the in vivo engraftment capability and broaden the scope of the T cell gene therapy. Engineering alternate immune cell subsets, such as CD4⁺ helper T cells, CD4⁺CD25⁺ regulatory T cells and $\gamma\delta$ T cells to produce specialized antigen-specific T cells is a current focus of research. Though, it is also important to minimize the safety concerns accompanying the in vivo use of genetically-engineered T cells, while pursuing such therapeutic approaches.

References

1. Parish CR. Cancer immunotherapy: the past, the present and the future. *Immunol Cell Biol* 2003; 81:106.
2. Finn OJ. Cancer vaccines: between the idea and the reality. *Nat Rev Immunol* 2003; 3:630.
3. Rosenberg SA, Yang JC, Restifo NP. Cancer immunotherapy: moving beyond current vaccines. *Nat Med* 2004; 10:909.
4. McKee MD, Fichera A, Nishimura MI. T cell immunotherapy. *Front Biosci* 2007; 12:919.
5. Marijt WA, Heemskerk MH, Kloosterboer FM, Goulmy E, Kester MG, van der Hoorn MA, et al. Hematopoiesis-restricted minor histocompatibility antigens HA-1- or HA-2-specific T cells can induce complete remissions of relapsed leukemia. *Proc Natl Acad Sci USA* 2003; 100:2742.
6. Akatsuka Y, Morishima Y, Kuzushima K, Kodera Y, Takahashi T. Minor histocompatibility antigens as targets for immunotherapy using allogeneic immune reactions. *Cancer Sci* 2007; 98:1139.
7. Kolb HJ, Mittermüller J, Clemm C, Holler E, Ledderose G, Brehm G, et al. Donor leukocyte transfusions for treatment of recurrent chronic myelogenous leukemia in marrow transplant patients. *Blood* 1990; 76:2462.
8. Kolb HJ, Schattenberg A, Goldman JM, Hertenstein B, Jacobsen N, Arcese W, et al. Graft-versus-leukemia effect of donor lymphocyte transfusions in marrow grafted patients. *Blood* 1995; 86:2041.

9. Bollard CM, Kuehnle I, Leen A, Rooney CM, Heslop HE. Adoptive immunotherapy for posttransplantation viral infections. *Biol Blood Marrow Transplant* 2004; 10:143.
10. Walter EA, Greenberg PD, Gilbert MJ, Finch RJ, Watanabe KS, Thomas ED, Riddell SR. Reconstitution of cellular immunity against cytomegalovirus in recipients of allogeneic bone marrow by transfer of T-cell clones from the donor. *N Engl J Med* 1995; 333:1038.
11. Einsele H, Roosnek E, Rufer N, Sinzger C, Riegler S, Löffler J, et al. Infusion of cytomegalovirus (CMV)-specific T cells for the treatment of CMV infection not responding to antiviral chemotherapy. *Blood* 2002; 99:3916.
12. Moss P, Rickinson A. Cellular immunotherapy for viral infection after HSC transplantation. *Nat Rev Immunol* 2005; 5:9.
13. Heslop HE, Ng CY, Li C, Smith CA, Loftin SK, Krance RA, et al. Long-term restoration of immunity against Epstein-Barr virus infection by adoptive transfer of gene-modified virus-specific T lymphocytes. *Nat Med* 1996; 2:551.
14. Rosenberg SA, Restifo NP, Yang JC, Morgan RA, Dudley ME. Adoptive cell transfer: a clinical path to effective cancer immunotherapy. *Nat Rev Cancer* 2008; 8:299.
15. Dudley ME, Rosenberg SA. Adoptive-cell-transfer therapy for the treatment of patients with cancer. *Nat Rev Cancer* 2003; 3:666.
16. Lotze MT, Grimm EA, Mazumder A, Strausser JL, Rosenberg SA. Lysis of fresh and cultured autologous tumor by human lymphocytes cultured in T-cell growth factor. *Cancer Res* 1981; 41:4420.
17. Rosenberg SA, Lotze MT, Muul LM, Chang AE, Avis FP, Leitman S, et al. A progress report on the treatment of 157 patients with advanced cancer using lymphokine-activated killer cells and interleukin-2 or high-dose interleukin-2 alone. *N Engl J Med* 1987; 316:889.
18. Rosenberg SA, Lotze MT, Yang JC, Topalian SL, Chang AE, Schwartzentruber DJ, et al. Prospective randomized trial of high-dose interleukin-2 alone or in conjunction with lymphokine-activated killer cells for the treatment of patients with advanced cancer. *J Natl Cancer Inst* 1993; 85:622.
19. Rosenberg SA, Yannelli JR, Yang JC, Topalian SL, Schwartzentruber DJ, Weber JS, et al. Treatment of patients with metastatic melanoma with autologous tumor-infiltrating lymphocytes and interleukin 2. *J Natl Cancer Inst* 1994; 86:1159.
20. Dudley ME, Wunderlich J, Nishimura MI, Yu D, Yang JC, Topalian SL, et al. Adoptive transfer of cloned melanoma-reactive T lymphocytes for the treatment of patients with metastatic melanoma. *J Immunother* 2001; 24:363.
21. Yee C, Thompson JA, Byrd D, Riddell SR, Roche P, Celis E, et al. Adoptive T cell therapy using antigen-specific CD8⁺ T cell clones for the treatment of patients with metastatic melanoma: in vivo persistence, migration and antitumor effect of transferred T cells. *Proc Natl Acad Sci USA* 2002; 99:16168.
22. Dudley ME, Wunderlich JR, Robbins PF, Yang JC, Hwu P, Schwartzentruber DJ, et al. Cancer regression and autoimmunity in patients after clonal repopulation with antitumor lymphocytes. *Science* 2002; 298:850.
23. Dudley ME, Wunderlich JR, Yang JC, Sherry RM, Topalian SL, Restifo NP, et al. Adoptive cell transfer therapy following non-myceloablative but lymphodepleting chemotherapy for the treatment of patients with refractory metastatic melanoma. *J Clin Oncol* 2005; 23:2346.
24. Dudley ME, Yang JC, Sherry R, Hughes MS, Royal R, Kammula U, et al. Adoptive cell therapy for patients with metastatic melanoma: evaluation of intensive myeloablative chemoradiation preparative regimens. *J Clin Oncol* 2008; 26:5233.
25. Levine BL, Humeau LM, Boyer J, MacGregor RR, Rebello T, Lu X, et al. Gene transfer in humans using a conditionally replicating lentiviral vector. *Proc Natl Acad Sci USA* 2006; 103:17372.
26. Mitsuysu RT, Anton PA, Deeks SG, Scadden DT, Connick E, Downs MT, et al. Prolonged survival and tissue trafficking following adoptive transfer of CD4zeta gene-modified autologous CD4(+) and CD8(+) T cells in human immunodeficiency virus-infected subjects. *Blood* 2000; 96:785.
27. Coccors M, Straetmans T, Govers C, Lamers C, Sleijfer S, Debets R. T cell receptor (TCR) gene therapy to treat melanoma: lessons from clinical and preclinical studies. *Expert Opin Biol Ther* 2010; 10:547.
28. Dembic Z, Haas W, Weiss S, McCubrey J, Kiefer H, von Boehmer H, et al. Transfer of specificity by murine alpha and beta T-cell receptor genes. *Nature* 1986; 320:232.
29. Cole DJ, Weil DP, Shamamian P, Rivoltini L, Kawakami Y, Topalian S, et al. Identification of MART-1-specific T-cell receptors: T cells utilizing distinct T-cell receptor variable and joining regions recognize the same tumor epitope. *Cancer Res* 1994; 54:5265.
30. Clay TM, Custer MC, Sachs J, Hwu P, Rosenberg SA, Nishimura MI. Efficient transfer of a tumor antigen-reactive TCR to human peripheral blood lymphocytes confers anti-tumor reactivity. *J Immunol* 1999; 163:507.
31. Clay TM, Custer MC, Spiess PJ, Nishimura MI. Potential use of T cell receptor genes to modify hematopoietic stem cells for the gene therapy of cancer. *Pathol Oncol Res* 1999; 5:3.
32. Calogero A, Hospers GA, Kruse KM, Schrier PI, Mulder NH, Hooijberg E, et al. Retargeting of a T cell line by anti MAGE-3/HLA-A2alpha beta TCR gene transfer. *Anticancer Res* 2000; 20:1793.
33. Roszkowski JJ, Yu DC, Rubinstein MR, McKee MD, Cole DJ, Nishimura MI. CD8-independent tumor cell recognition is a property of the T cell receptor and not the T cell. *J Immunol* 2003; 170:2582.
34. Morgan RA, Dudley ME, Yu YY, Zheng Z, Robbins PF, Theoret MR, et al. High efficiency TCR gene transfer into primary human lymphocytes affords avid recognition of melanoma tumor antigen glycoprotein 100 and does not alter the recognition of autologous melanoma antigens. *J Immunol* 2003; 171:3287.
35. Stanislawski T, Voss RH, Lotz C, Sadovnikova E, Willemsen RA, Kuball J, et al. Circumventing tolerance to a human MDM2-derived tumor antigen by TCR gene transfer. *Nat Immunol* 2001; 2:962.
36. Zhang Y, Liu Y, Moxley KM, Golden-Mason L, Hughes MG, Liu T, et al. Transduction of human T cells with a novel T-cell receptor confers anti-HCV reactivity. *PLoS Pathog* 6:1001018.
37. Orentas RJ, Roskopf SJ, Nolan GP, Nishimura MI. Retroviral transduction of a T cell receptor specific for an Epstein-Barr virus-encoded peptide. *Clin Immunol* 2001; 98:220.
38. Norell H, Zhang Y, McCracken J, Martins da Palma T, Leshner A, Liu Y, et al. CD34-based enrichment of genetically engineered human T cells for clinical use results in dramatically enhanced tumor targeting. *Cancer Immunol Immunother* 59:851.
39. McKee MD, Roszkowski JJ, Nishimura MI. T cell avidity and tumor recognition: implications and therapeutic strategies. *J Transl Med* 2005; 3:35.
40. Benihoud K, Yeh P, Perricaudet M. Adenovirus vectors for gene delivery. *Curr Opin Biotechnol* 1999; 10:440.
41. Weber E, Anderson WF, Kasahara N. Recent advances in retrovirus vector-mediated gene therapy: teaching an old vector new tricks. *Curr Opin Mol Ther* 2001; 3:439.
42. Cavalieri S, Cazzaniga S, Geuna M, Magnani Z, Bordignon C, Naldini L, et al. Human T lymphocytes transduced by lentiviral vectors in the absence of TCR activation maintain an intact immune competence. *Blood* 2003; 102:497.
43. Zhou X, Cui Y, Huang X, Yu Z, Thomas AM, Ye Z, et al. Lentivirus-mediated gene transfer and expression in established human tumor antigen-specific cytotoxic T cells and primary unstimulated T cells. *Hum Gene Ther* 2003; 14:1089.
44. Cohen CJ, Zhao Y, Zheng Z, Rosenberg SA, Morgan RA. Enhanced antitumor activity of murine-human hybrid T-cell receptor (TCR) in human lymphocytes is associated with improved pairing and TCR/CD3 stability. *Cancer Res* 2006; 66:8878.
45. Cohen CJ, Li YF, El-Gamil M, Robbins PF, Rosenberg SA, Morgan RA. Enhanced antitumor activity of T cells engineered to express T-cell receptors with a second disulfide bond. *Cancer Res* 2007; 67:3898.
46. Li Y, Moysey R, Molloy PE, Vuidepot AL, Mahon T, Baston E, et al. Directed evolution of human T-cell receptors with picomolar affinities by phage display. *Nat Biotechnol* 2005; 23:349.
47. Theobald M, Biggs J, Dittmer D, Levine AJ, Sherman LA. Targeting p53 as a general tumor antigen. *Proc Natl Acad Sci USA* 1995; 92:11993.
48. Morgan RA, Dudley ME, Wunderlich JR, Hughes MS, Yang JC, Sherry RM, et al. Cancer regression in patients after transfer of genetically engineered lymphocytes. *Science* 2006; 314:126.
49. Johnson LA, Morgan RA, Dudley ME, Cassard L, Yang JC, Hughes MS, et al. Gene therapy with human and mouse T-cell receptors mediates cancer regression and targets normal tissues expressing cognate antigen. *Blood* 2009; 114:535.
50. Parkhurst MR, Yang JC, Langan RC, Dudley ME, Nathan DA, Feldman SA, et al. T Cells Targeting Carcinoembryonic Antigen Can Mediate Regression of Metastatic Colorectal Cancer but Induce Severe Transient Colitis. *Mol Ther*.
51. Gross G, Waks T, Eshhar Z. Expression of immunoglobulin-T-cell receptor chimeric molecules as functional receptors with antibody-type specificity. *Proc Natl Acad Sci USA* 1989; 86:10024.
52. Gross G, Gorochoff G, Waks T, Eshhar Z. Generation of effector T cells expressing chimeric T cell receptor with antibody type-specificity. *Transplant Proc* 1989; 21:127.
53. Turatti F, Figini M, Balladore E, Alberti P, Casalini P, Marks JD, et al. Redirected activity of human antitumor chimeric immune receptors is governed by antigen and receptor expression levels and affinity of interaction. *J Immunother* 2007; 30:684.
54. Lamers CH, Sleijfer S, Vulto AG, Kruit WH, Kliffen M, Debets R, et al. Treatment of metastatic renal cell carcinoma with autologous T-lymphocytes genetically retargeted against carbonic anhydrase IX: first clinical experience. *J Clin Oncol* 2006; 24:20.
55. Haynes NM, Trapani JA, Teng MW, Jackson JT, Cerruti L, Jane SM, et al. Rejection of syngeneic colon carcinoma by CTLs expressing single-chain antibody receptors codelivering CD28 costimulation. *J Immunol* 2002; 169:5780.
56. Westwood JA, Smyth MJ, Teng MW, Moeller M, Trapani JA, Scott AM, et al. Adoptive transfer of T cells modified with a humanized chimeric receptor gene inhibits growth of Lewis-Y-expressing tumors in mice. *Proc Natl Acad Sci USA* 2005; 102:19051.
57. Daly T, Royal RE, Kershaw MH, Treisman J, Wang G, Li W, et al. Recognition of human colon cancer by T cells transduced with a chimeric receptor gene. *Cancer Gene Ther* 2000; 7:284.
58. Ma Q, Safar M, Holmes E, Wang Y, Boynton AL, Junghans RP. Anti-prostate specific membrane antigen designer T cells for prostate cancer therapy. *Prostate* 2004; 61:12.
59. Yun CO, Nolan KF, Beecham EJ, Reisfeld RA, Junghans RP. Targeting of T lymphocytes to melanoma cells through chimeric anti-GD3 immunoglobulin T-cell receptors. *Neoplasia* 2000; 2:449.

60. Cooper LJ, Al-Kadhimi Z, Serrano LM, Pfeiffer T, Olivares S, Castro A, et al. Enhanced antilymphoma efficacy of CD19-redirected influenza MP1-specific CTLs by cotransfer of T cells modified to present influenza MP1. *Blood* 105:1622.
61. Finney HM, Lawson AD, Bebbington CR, Weir AN. 1998. Chimeric receptors providing both primary and costimulatory signaling in T cells from a single gene product. *J Immunol* 2005; 161:2791.
62. Maher J, Brentjens RJ, Gunset G, Riviere I, Sadelain M. Human T-lymphocyte cytotoxicity and proliferation directed by a single chimeric TCRzeta/CD28 receptor. *Nat Biotechnol* 2002; 20:70.
63. Finney HM, Akbar AN, Lawson AD. Activation of resting human primary T cells with chimeric receptors: costimulation from CD28, inducible costimulator, CD134 and CD137 in series with signals from the TCR zeta chain. *J Immunol* 2004; 172:104.
64. Zhong XS, Matsushita M, Plotkin J, Riviere I, Sadelain M. Chimeric antigen receptors combining 4-1BB and CD28 signaling domains augment PI3kinase/AKT/Bcl-XL activation and CD8⁺ T cell-mediated tumor eradication. *Mol Ther* 18:413.
65. Carpenito C, Milone MC, Hassan R, Simonet JC, Lakhal M, Suhoski MM, et al. Control of large, established tumor xenografts with genetically retargeted human T cells containing CD28 and CD137 domains. *Proc Natl Acad Sci USA* 2009; 106:3360.
66. Till BG, Jensen MC, Wang J, Chen EY, Wood BL, Greisman HA, et al. Adoptive immunotherapy for indolent non-Hodgkin lymphoma and mantle cell lymphoma using genetically modified autologous CD20-specific T cells. *Blood* 2008; 112:2261.
67. Kershaw MH, Westwood JA, Parker LL, Wang G, Eshhar Z, Mavroukakis SA, et al. A phase I study on adoptive immunotherapy using gene-modified T cells for ovarian cancer. *Clin Cancer Res* 2006; 12:6106.
68. Pule MA, Savoldo B, Myers GD, Rossig C, Russell HV, Dotti G, et al. Virus-specific T cells engineered to coexpress tumor-specific receptors: persistence and antitumor activity in individuals with neuroblastoma. *Nat Med* 2008; 14:1264.
69. Park JR, Digiusto DL, Slovak M, Wright C, Naranjo A, Wagner J, et al. Adoptive transfer of chimeric antigen receptor re-directed cytolytic T lymphocyte clones in patients with neuroblastoma. *Mol Ther* 2007; 15:825.
70. Morgan RA, Yang JC, Kitano M, Dudley ME, Laurencot CM, Rosenberg SA. Case report of a serious adverse event following the administration of T cells transduced with a chimeric antigen receptor recognizing ERBB2. *Mol Ther* 18:843.
71. Brentjens R, Yeh R, Bernal Y, Riviere I, Sadelain M. Treatment of chronic lymphocytic leukemia with genetically targeted autologous T cells: case report of an unforeseen adverse event in a phase I clinical trial. *Mol Ther* 18:666.